

**DEMO**

**100%  
Money  
Back  
Guarantee**

**Instant  
Download  
After  
Purchase**

**PDF**  
FILE FORMAT

**90  
Days  
Free  
Updates**

**American Academy of  
Cosmetic Dentistry**

# **AACD-AACD**

## **Q&As**

2020 Latest EXAMSDUMPS AACD PDF Dumps Download. Following Questions and Answers are all new published by AACD Official Exam Center

**Latest AACD-AACD Dumps**

**AACD-AACD Practice Test**

**AACD-AACD Study Guide**

**Pass AACD AACD Exam with  
100% Guarantee**

Free Download Real  
Questions & Answers  
PDF and VCE file from:



+13069525559



<https://examsdumps.co/>



<https://www.facebook.com/examsdumps.co/>

**AACD**

**American Academy of Cosmetic Dentistry**

**Exam A****QUESTION 1**

Which of the following would be the main advantage of utilizing a layering technique with composite resin in restoring a disto-occlusal Class II restoration?

- A. Higher degree of polymerization and lower degree of tooth sensitivity
- B. Higher shear strength and wear resistance
- C. Higher tensile strength and modulus of elasticity
- D. Better compressive strength

**Correct Answer:** A

**Section:** (none)

**Explanation**

**Explanation/Reference:**

Explanation:

**QUESTION 2**

Which of the following is correct concerning tooth preparation for diastema closure with bonded porcelain restorations?

- A. Keep all margins supragingival or equigingival
- B. Full crown preparations are best used to close spaces
- C. Subgingival interdental margins will allow for improved emergence profiles
- D. Mini-wings must always be used to close a black triangle interdentally

**Correct Answer:** C

**Section:** (none)

**Explanation**

**Explanation/Reference:**

References:

**QUESTION 3**

What is the BEST property of condensible composites?

- A. Lower filler content to allow lateral and vertical flow ensuring intimate contact
- B. Less stickiness and stiffer consistency and have a wear ratio similar to amalgam
- C. A filler loading of fibrous, porous, or irregular particles, and a filler percentage of 40% to 50% filled
- D. Clinical evidence of strength and polymerization shrinkage superior to conventional composites

**Correct Answer:** D

**Section:** (none)

**Explanation**

**Explanation/Reference:**

Explanation:

**QUESTION 4**

A white opacity near the incisal one-third of a permanent incisor is MOST likely the result of what?

- A. Ingestion of excessive fluoride by the patient's mother during pregnancy
- B. Ingestion of excessive fluoride by the patient at the age of 1 - 3 years
- C. Antibiotic therapy by mother during pregnancy
- D. Trauma or infection of the patient's deciduous incisor

**Correct Answer:** D

**Section:** (none)

**Explanation**

**Explanation/Reference:**

Explanation:

**QUESTION 5**

When preparing for posterior porcelain to zirconia crown, the recommended occlusal reduction should be:

- A. 1.0 mm.
- B. 1.5 mm.
- C. 2.0 mm.
- D. 2.5 mm.

**Correct Answer:** B

**Section:** (none)

**Explanation**

**Explanation/Reference:**

References:

**BUY NOW**